

27. Godoy D, Randle G, Simpson AJ, Aanensen DM, Pitt TL, Kinoshita R, et al. Multilocus sequence typing and evolutionary relationships among the causative agents of melioidosis and glanders, *Burkholderia pseudomallei* and *Burkholderia mallei*. *J Clin Microbiol*. 2003;41:2068–79. <http://dx.doi.org/10.1128/JCM.41.5.2068-2079.2003>
28. Brett PJ, Deshazer D, Woods DE. Characterization of *Burkholderia pseudomallei* and *Burkholderia pseudomallei*-like strains. *Epidemiol Infect*. 1997;118:137–48. <http://dx.doi.org/10.1017/S095026889600739X>
29. Bremmelgaard A, Bygbjerg I, Hoiby N. Microbiological and immunological studies in a case of human melioidosis diagnosed in Denmark. *Scand J Infect Dis*. 1982;14:271–5.
30. Wall RA, Mabey DC, Corrah PT, Peters L. A case of melioidosis in West Africa. *J Infect Dis*. 1985;152:424–5. <http://dx.doi.org/10.1093/infdis/152.2.424a>
31. Issack MI, Bundhun CD, Gokhool H. Melioidosis in Mauritius. *Emerg Infect Dis*. 2005;11:139–40. <http://dx.doi.org/10.3201/eid1101.040605>
32. Martinet O, Pac Soo AM, Knezynski M. Melioidosis: regarding a case acquired in Madagascar and two nosocomial cases [in French]. *Bull Soc Pathol Exot*. 2004;97:366–70.
33. Borgherini G, Poubeau P, Paganin F, Picot S, Michault A, Thibault F, et al. Melioidosis: an imported case from Madagascar. *J Travel Med*. 2006;13:318–20. <http://dx.doi.org/10.1111/j.1708-8305.2006.00050.x>
34. Amezyane T, Lecoules S, Algayres JP. Mycotic iliac aneurysm associated with *Burkholderia pseudomallei*. *Int J Infect Dis*. 2010;14(Suppl 3):e381–2. <http://dx.doi.org/10.1016/j.ijid.2009.07.008>
35. Bertherat E, Thullier P, Shako JC, England K, Koné ML, Arntzen L, et al. Lessons learned about pneumonic plague diagnosis from two outbreaks, Democratic Republic of the Congo. *Emerg Infect Dis*. 2011;17:778–84. <http://dx.doi.org/10.3201/eid1705.100029>
36. Rattanavong S, Wuthiekanun V, Langla S, Amornchai P, Sirisouk J, Phetsouvanh R, et al. Randomized soil survey of the distribution of *Burkholderia pseudomallei* in rice fields in Laos. *Appl Environ Microbiol*. 2011;77:532–6. <http://dx.doi.org/10.1128/AEM.01822-10>
37. Kaestli M, Mayo M, Harrington G, Ward L, Watt F, Hill JV, et al. Landscape changes influence the occurrence of the melioidosis bacterium *Burkholderia pseudomallei* in soil in northern Australia. *PLoS Negl Trop Dis*. 2009;3:e364. <http://dx.doi.org/10.1371/journal.pntd.0000364>
38. Ussery DW, Kiil K, Lagesen K, Sicheritz-Ponten T, Bohlin J, Wassenaar TM. The genus *Burkholderia*: analysis of 56 genomic sequences. *Genome Dyn*. 2009;6:140–57. <http://dx.doi.org/10.1159/000235768>
39. Hantrakun V, Rongkard P, Amonchi P, Sarunporn T, Langla S, Wuthiekanun V, et al. Presence of environmental *B. pseudomallei*, *B. thailandensis* and putative *B. thailandensis* with *B. ps*-like CPS variant in east Thailand. Abstract in: Proceedings of the 7th World Melioidosis Congress; 2013 Sep 18–20; Bangkok, Thailand.
40. Peacock SJ, Cheng AC, Currie BJ, Dance DA. The use of positive serological tests as evidence of exposure to *Burkholderia pseudomallei*. *Am J Trop Med Hyg*. 2011;84:1021–2. <http://dx.doi.org/10.4269/ajtmh.2011.11-0114a>

Address for correspondence: W. Joost Wiersinga, Center for Infection and Immunity Amsterdam, Dept. of Internal Medicine, Academic Medical Center, University of Amsterdam, Meibergdreef 9, Room G2-132, 1105 AZ Amsterdam, the Netherlands; email: w.j.wiersinga@amc.uva.nl

etymologia

Glanders [glan'dærz]

From the Old French *glandres* (“glands”) describing the enlargement of the parotid or submaxillary lymph nodes that is pathognomonic of the disease, glanders is a contagious disease of horses. Glanders is caused by *Burkholderia mallei* and is communicable to humans but should not be confused with human melioidosis, caused by *Burkholderia pseudomallei*. The chronic, cutaneous form of glanders presents as ulcerated skin lesions along major lymph and blood vessels and is known as farcy (from the Latin *farcire*, “sausage”). Among the first descriptions of glanders is in

the writings of Aristotle: “The ass suffers chiefly from one particular disease which they call ‘melis.’” In later writings, “melis” became “malleus,” which became a generic term for epizootics. Glanders has been eliminated in many industrialized countries, including in the United States, where there have been no naturally acquired human or animal cases since World War II. More recently, glanders has been reemerging in parts of the world; since 2000, outbreaks in horses have been reported in North Africa, the Middle East, and other areas.

Sources

1. Dorland’s Illustrated Medical Dictionary. 32nd ed. Philadelphia: Elsevier Saunders; 2012.
2. Glanders. Centers for Disease Control and Prevention [cited 2014 Sep 25]. <http://www.cdc.gov/glanders/index.html>.
3. Wilkinson L. Glanders: medicine and veterinary medicine in common pursuit of a contagious disease. *Med Hist*. 1981;25:363–84. <http://dx.doi.org/10.1017/S0025727300034876>

Address for correspondence: Ronnie Henry, Centers for Disease Control and Prevention, 1600 Clifton Rd NE, Mailstop E03, Atlanta, GA 30329-4027, USA; email: boq3@cdc.gov

DOI: <http://dx.doi.org/10.3201/eid2101.ET2101>