


## 3-*N*-(2-[<sup>18</sup>F]Fluoroethyl)piperone [<sup>18</sup>F]FESP

Kam Leung, PhD<sup>1</sup>

Created: October 7, 2005; Updated: January 18, 2012.

|                                     |                                                                                                                                |
|-------------------------------------|--------------------------------------------------------------------------------------------------------------------------------|
| <b>Chemical name:</b>               | 3- <i>N</i> -(2-[ <sup>18</sup> F]Fluoroethyl)piperone                                                                         |
| <b>Abbreviated name:</b>            | [ <sup>18</sup> F]FESP                                                                                                         |
| <b>Synonym:</b>                     |                                                                                                                                |
| <b>Agent Category:</b>              | Compound                                                                                                                       |
| <b>Target:</b>                      | D <sub>2</sub> dopamine receptor and 5-HT <sub>2</sub> serotonin receptor                                                      |
| <b>Target Category:</b>             | Receptor                                                                                                                       |
| <b>Method of detection:</b>         | PET                                                                                                                            |
| <b>Source of signal / contrast:</b> | <sup>18</sup> F                                                                                                                |
| <b>Activation:</b>                  | No                                                                                                                             |
| <b>Studies:</b>                     | <ul style="list-style-type: none"> <li><i>In vitro</i></li> <li>Rodents</li> <li>Non-human primates</li> <li>Humans</li> </ul> |



Click on the above structure for additional information in [PubChem](#).

## Background

[[PubMed](#)]

Dopamine, a neurotransmitter, plays an important role in the mediation of movement, cognition, and emotion (1, 2). Dopamine shortage plays a role in various neuropsychiatric disorders, such as Parkinson's disease (PD), schizophrenia, autism, attention deficit hyperactivity disorder, and drug abuse. Two subtypes of dopamine receptors, D<sub>1</sub> and D<sub>2</sub>, were well-characterized pharmacologically and biochemically (3). D<sub>2</sub> dopamine receptors have been implicated in the pathophysiology of PD, Alzheimer's disease, and schizophrenia (4).

Serotonin (5-hydroxytryptamine, 5-HT) has diverse physiological roles as a neurotransmitter in the central nervous system (5). It also is a regulator of smooth muscle function and platelet aggregation. The brain cortical

5-HT system has been implicated in several neuropsychiatric disorders, including major depression, anxiety, obsessive-compulsive disorder, and schizophrenia (6, 7).

Spiperone and its analog, 3-*N*-(2-fluoroethyl)spiperone (FESP) are high-affinity D<sub>2</sub> dopamine and 5-HT<sub>2</sub> serotonin receptor antagonists, showing a low affinity for  $\alpha$ 1-adrenergic receptors and marginal binding to other receptors (8). 3-*N*-(2-[<sup>18</sup>F]Fluoroethyl)spiperone ([<sup>18</sup>F]FESP) has been studied as a positron emission tomography (PET) tracer for imaging D<sub>2</sub> and 5HT<sub>2</sub> receptor densities. It has also used as a reporter probe for dopamine D<sub>2</sub> receptor in imaging transgene expression in rodents (9-11).

## Related Resource Links:

- Chapters in MICAD ([5-HT<sub>2A</sub>](#), [Dopamine receptors](#))
- Gene information in NCBI ([5-HT<sub>2A</sub>](#), [D<sub>2</sub> receptor](#), [D<sub>3</sub> receptor](#))
- Articles in Online Mendelian Inheritance in Man (OMIM) ([5-HT<sub>2A</sub>](#), [D<sub>2</sub> receptor](#), [D<sub>3</sub> receptor](#))
- Clinical trials ([5-HT<sub>2A</sub>](#), [Dopamine receptors](#))
- Drug information in Food and Drug Administration ([5-HT<sub>2A</sub>](#), [Dopamine receptors](#))

## Synthesis

[PubMed]

[<sup>18</sup>F]FESP can be synthesized by either N-alkylation or nucleophilic substitution (12-15). Alkylation of the amide nitrogen in spiperone by [<sup>18</sup>F]fluorobromoethane in the presence of NaOH provided [<sup>18</sup>F]FESP in 35-40% radiochemical yield (5-8% overall radiochemical yield). [<sup>18</sup>F]Fluorobromoethane was easily prepared by the reaction of 1,2-bromoethane with potassium [<sup>18</sup>F]fluoride Kryptofix complex in 15-20% radiochemical yield. The radiochemical purity of [<sup>18</sup>F]FESP was found to be >99% by high-performance liquid chromatography (HPLC). The specific activity was 185-370 GBq/ $\mu$ mol (5-10 Ci/ $\mu$ mol) at the end of synthesis with a total synthesis time of 70 min.

A one-step, one-pot synthesis of [<sup>18</sup>F]FESP using nucleophilic fluorination of 3-(2'-bromoethyl)spiperone or 3-(2'-methylsulfonyloxyethyl)spiperone with K[<sup>18</sup>F]F/Kryptofix2.2.2 in acetonitrile in <60 min was performed (12). This method provided a 99% chemical purity and a specific activity of 74-370 GBq/ $\mu$ mol (2-10 Ci/ $\mu$ mol) at the end of synthesis (EOS). The radiochemical yield was 30-40% (EOS).

## In Vitro Studies: Testing in Cells and Tissues

[PubMed]

FESP was reported to have selective binding affinity to D<sub>2</sub> (striatum) and 5-HT<sub>2</sub> (frontal cortex) receptor sites in homogenates of rat brain membranes (8). The *K<sub>i</sub>* values for D<sub>2</sub> and 5-HT<sub>2</sub> were 0.44 nM and 0.57 nM, respectively. It has a *K<sub>i</sub>* value of 23 nM for the  $\alpha$ 1-adrenergic receptor in rat forebrain membrane and negligible binding to other sites.

## Animal Studies

### Rodents

[PubMed]

Biodistribution studies in rats showed a high uptake of radioactivity in the lung (1.80% injected dose (ID)/g), kidney (1.00% ID/g), spleen (0.97% ID/g), and liver (0.82% ID/g) at 15 min after injection of [<sup>18</sup>F]FESP (16). The brain had a low uptake of 0.18% ID/g. Low radioactivity was observed in bone tissue, indicating little

defluorination. [<sup>18</sup>F]FESP remained 90% intact in the brain striatum, 32% in the cerebellum, and 46% in the frontal cortex for up to 4 h after injection. In mice, [<sup>18</sup>F]FESP remained 90% intact in the brain striatum, and 62% in the cerebellum for up to 3.5 h after injection (17). [<sup>18</sup>F]FESP was metabolized peripherally with only 11% intact [<sup>18</sup>F]FESP in rat plasma at 2 h after injection (16).

## Other Non-Primate Mammals

[PubMed]

No publication is currently available.

## Non-Human Primates

[PubMed]

[<sup>18</sup>F]FESP PET studies in non-human primates have provided useful assessment of the D<sub>2</sub> and 5-HT<sub>2</sub> receptor in the brain (16-21). Coenen et al. (18) showed a selective uptake in striatum (0.048% ID/cm<sup>3</sup>) over the frontal cortex (0.0088% ID/cm<sup>3</sup>) and in the cerebellum (0.0056% ID/cm<sup>3</sup>) in baboon brains at 4 h after injection. The striatum D<sub>2</sub> receptor sites in the striatum were saturable. The striatum [<sup>18</sup>F]FESP uptake was blocked by pretreatment with excess unlabeled FESP (0.5 mg/kg) and (+)-butaclamol (0.5 mg/kg), a D<sub>1</sub>/D<sub>2</sub> antagonist (22). Ritanserin and ketanserin, 5-HT<sub>2</sub> antagonists, had no effect on [<sup>18</sup>F]FESP uptake in the striatum and blocked only 15-20% [<sup>18</sup>F]FESP uptake in the frontal cortex. [<sup>18</sup>F]FESP arterial plasma activity rapidly decreased to 25% at 1 h and 8% at 4 h after injection. D<sub>2</sub> receptor density ( $B_{\max}$ ) was estimated by two PET studies to be  $25.9 \pm 12.7$  pmol/g (19) and  $37.5 \pm 11.5$  pmol/g (21). Using serial PET scans, 5-HT<sub>2</sub> receptor density was estimated to be  $35.6 \pm 10.9$  pmol/g in the baboon frontal cortex (20).

## Human Studies

[[PubMed]]

Human dosimetry was estimated based on human dynamic PET scans after injection of 37 MBq (1 mCi) [<sup>18</sup>F]FESP (23). The gallbladder received the highest dose (0.207 mGy/MBq or 797 mrad/mCi). Other organs that received high doses were the urinary bladder (0.083 mGy/MBq or 308 mrad/mCi) and liver (0.065 mGy/MBq or 110 mrad/mCi). The total body dose was 0.0119 mGy/MBq (44 mrad/mCi).

[<sup>18</sup>F]FESP PET studies of D<sub>2</sub> receptor distribution in human brain were reported, showing a localization of radioactivity in the striatum (16, 24, 25). Only about 1% of [<sup>18</sup>F]FESP entered the brain. [<sup>18</sup>F]FESP was metabolized peripherally with 54% intact [<sup>18</sup>F]FESP in arterial plasma at 2 h after injection (16). Striatal-to-cerebellum ratio and kinetic constants are commonly used as analytical parameters in [<sup>18</sup>F]FESP PET studies. Wienhard et al. (25) reported on [<sup>18</sup>F]FESP PET studies in 30 patients with various disorders related to the dopaminergic system and in 6 normal subjects. PET brain scans of normal subjects showed that the highest uptake was in the caudate and pituitary, followed by the cortex and cerebellum at 3 h after injection of 185 MBq (5 mCi) [<sup>18</sup>F]FESP. PET scans of patients showed decreases in the number of available D<sub>2</sub> receptors in the striatum and striatum-to-cerebellum ratio. Patients with brain damage also showed both low [<sup>18</sup>F]FDG and [<sup>18</sup>F]FESP uptake in the striatum.

5-HT<sub>2</sub> receptor distribution in human brain were studied by [<sup>18</sup>F]FESP PET in 20 healthy volunteers and 34 patients with unipolar depression (26). There was a significant reduction of binding capacity in 19 non-drug-treated patients as compared with healthy controls in the cortical regions but not in the striatum. In paroxetine-treated patients ( $n = 15$ ), there was no significant difference between the drug-treated patients and healthy volunteers. Paroxetine is a specific serotonin reuptake inhibitor. Paroxetine treatment led to remission in these patients. Chronic treatment with fluvoxamine (27), another specific serotonin reuptake inhibitor, significantly

increased the *in vivo* binding of [ $^{18}\text{F}$ ]FESP in the frontal and occipital cortex but not in the striatum of drug-naïve, unipolar, depressed patients ( $n = 9$ ).

[ $^{18}\text{F}$ ]FESP PET is useful for objective monitoring of  $\text{D}_2$  and  $5\text{-HT}_2$  receptor sites in patients with serotonergic and dopaminergic disorders. It revealed stage-related changes in depression and dopaminergic disorders.

[ $^{18}\text{F}$ ]FESP has also demonstrated its usefulness as a PET tracer in the imaging of non-functioning pituitary adenomas from meningiomas and craniopharyngiomas (28).

## References

1. Carbon M., Ghilardi M.F., Feigin A., Fukuda M., Silvestri G., Mentis M.J., Ghez C., Moeller J.R., Eidelberg D. *Learning networks in health and Parkinson's disease: reproducibility and treatment effects*. . Hum Brain Mapp. 2003;19(3):197–211. PubMed PMID: 12811735.
2. Chesselet M.F., Delfs J.M. *Basal ganglia and movement disorders: an update*. . Trends Neurosci. 1996;19(10):417–22. PubMed PMID: 8888518.
3. Stoof J.C., Kebabian J.W. *Two dopamine receptors: biochemistry, physiology and pharmacology*. . Life Sci. 1984;35(23):2281–96. PubMed PMID: 6390056.
4. Seeman P., Bzowej N.H., Guan H.C., Bergeron C., Reynolds G.P., Bird E.D., Riederer P., Jellinger K., Tourtellotte W.W. *Human brain D1 and D2 dopamine receptors in schizophrenia, Alzheimer's, Parkinson's, and Huntington's diseases*. . Neuropsychopharmacology. 1987;1(1):5–15. PubMed PMID: 2908095.
5. Lucki I. *The spectrum of behaviors influenced by serotonin*. . Biol Psychiatry. 1998;44(3):151–62. PubMed PMID: 9693387.
6. Abi-Dargham A., Laruelle M., Aghajanian G.K., Charney D., Krystal J. *The role of serotonin in the pathophysiology and treatment of schizophrenia*. . J Neuropsychiatry Clin Neurosci. 1997;9(1):1–17. PubMed PMID: 9017523.
7. Charney D.S. *Monoamine dysfunction and the pathophysiology and treatment of depression*. . J Clin Psychiatry. 1998;59 Suppl 14:11–4. PubMed PMID: 9818625.
8. Goffinet A.M., Leysen J., Labar D. *In vitro pharmacological profile of 3-N-(2-fluoroethyl)spiperone*. . J Cereb Blood Flow Metab. 1990;10(1):140–2. PubMed PMID: 2137131.
9. MacLaren D.C., Gambhir S.S., Satyamurthy N., Barrio J.R., Sharfstein S., Toyokuni T., Wu L., Berk A.J., Cherry S.R., Phelps M.E., Herschman H.R. *Repetitive, non-invasive imaging of the dopamine D2 receptor as a reporter gene in living animals*. . Gene Ther. 1999;6(5):785–91. PubMed PMID: 10505102.
10. Yaghoubi S.S., Wu L., Liang Q., Toyokuni T., Barrio J.R., Namavari M., Satyamurthy N., Phelps M.E., Herschman H.R., Gambhir S.S. *Direct correlation between positron emission tomographic images of two reporter genes delivered by two distinct adenoviral vectors*. . Gene Ther. 2001;8(14):1072–80. PubMed PMID: 11526454.
11. Chen I.Y., Wu J.C., Min J.J., Sundaresan G., Lewis X., Liang Q., Herschman H.R., Gambhir S.S. *Micro-positron emission tomography imaging of cardiac gene expression in rats using bicistronic adenoviral vector-mediated gene delivery*. . Circulation. 2004;109(11):1415–20. PubMed PMID: 15007006.
12. Satyamurthy N., Bida G.T., Barrio J.R., Luxen A., Mazziotta J.C., Huang S.C., Phelps M.E. *No-carrier-added 3-(2'-[ $^{18}\text{F}$ ]fluoroethyl)spiperone, a new dopamine receptor-binding tracer for positron emission tomography*. . Int J Rad Appl Instrum B. 1986;13(6):617–24. PubMed PMID: 3494003.
13. Satyamurthy N., Barrio J.R., Bida G.T., Huang S.C., Mazziotta J.C., Phelps M.E. *3-(2'-[ $^{18}\text{F}$ ]fluoroethyl)spiperone, a potent dopamine antagonist: synthesis, structural analysis and in-vivo utilization in humans*. . Int J Rad Appl Instrum [A]. 1990;41(2):113–29. PubMed PMID: 2158942.
14. Chi D.Y., Kilbourn M.R., Katzenellenbogen J.A., Brodack J.W., Welch M.J. *Synthesis of no-carrier-added N-([ $^{18}\text{F}$ ]fluoroalkyl)spiperone derivatives*. . Int J Rad Appl Instrum [A]. 1986;37(12):1173–80. PubMed PMID: 3028983.

15. Kiesewetter D.O., Eckelman W.C., Cohen R.M., Finn R.D., Larson S.M. *Syntheses and D2 receptor affinities of derivatives of spiperone containing aliphatic halogens.* . Int J Rad Appl Instrum [A]. 1986;37(12):1181–8. PubMed PMID: 3028984.
16. Barrio J.R., Satyamurthy N., Huang S.C., Keen R.E., Nissenson C.H., Hoffman J.M., Ackermann R.F., Bahn M.M., Mazziotta J.C., Phelps M.E. *3-(2'-[18F]fluoroethyl)spiperone: in vivo biochemical and kinetic characterization in rodents, nonhuman primates, and humans.* . J Cereb Blood Flow Metab. 1989;9(6):830–9. PubMed PMID: 2531146.
17. Coenen H.H., Laufer P., Stocklin G., Wienhard K., Pawlik G., Bocher-Schwarz H.G., Heiss W.D. *3-N-(2-[18F]-fluoroethyl)-spiperone: a novel ligand for cerebral dopamine receptor studies with pet.* . Life Sci. 1987;40(1):81–8. PubMed PMID: 2948091.
18. Coenen H.H., Wienhard K., Stocklin G., Laufer P., Hebold I., Pawlik G., Heiss W.D. *PET measurement of D2 and S2 receptor binding of 3-N-[(2'-18F)fluoroethyl]spiperone in baboon brain.* . Eur J Nucl Med. 1988;14(2):80–7. PubMed PMID: 2968911.
19. Huang S.C., Bahn M.M., Barrio J.R., Hoffman J.M., Satyamurthy N., Hawkins R.A., Mazziotta J.C., Phelps M.E. *A double-injection technique for in vivo measurement of dopamine D2-receptor density in monkeys with 3-(2'-[18F]fluoroethyl)spiperone and dynamic positron emission tomography.* . J Cereb Blood Flow Metab. 1989;9(6):850–8. PubMed PMID: 2531148.
20. Jovkar S., Wienhard K., Coenen H.H., Pawlik G., Heiss W.D. *Quantification of baboon cortical S2 serotonin receptors in vivo with 3-N-(2'-F18)fluoroethylspiperone and positron emission tomography.* . Eur J Nucl Med. 1991;18(3):158–63. PubMed PMID: 2040338.
21. Jovkar S., Wienhard K., Pawlik G., Coenen H.H. *The quantitative analysis of D2-dopamine receptors in baboon striatum in vivo with 3-N-[2'-18F]fluoroethylspiperone using positron emission tomography.* . J Cereb Blood Flow Metab. 1990;10(5):720–6. PubMed PMID: 2384543.
22. Schetz J.A., Kim O.J., Sibley D.R. *Pharmacological characterization of mammalian D1 and D2 dopamine receptors expressed in Drosophila Schneider-2 cells.* . J Recept Signal Transduct Res. 2003;23(1):99–109. PubMed PMID: 12680592.
23. Herzog H., Coenen H.H., Kuwert T., Langen K.J., Feinendegen L.E. *Quantification of the whole-body distribution of PET radiopharmaceuticals, applied to 3-N-[(18F]fluoroethyl)spiperone.* . Eur J Nucl Med. 1990;16(2):77–83. PubMed PMID: 2311622.
24. Bahn M.M., Huang S.C., Hawkins R.A., Satyamurthy N., Hoffman J.M., Barrio J.R., Mazziotta J.C., Phelps M.E. *Models for in vivo kinetic interactions of dopamine D2-neuroreceptors and 3-(2'-[18F]fluoroethyl)spiperone examined with positron emission tomography.* . J Cereb Blood Flow Metab. 1989;9(6):840–9. PubMed PMID: 2531147.
25. Wienhard K., Coenen H.H., Pawlik G., Rudolf J., Laufer P., Jovkar S., Stocklin G., Heiss W.D. *PET studies of dopamine receptor distribution using [18F]fluoroethylspiperone: findings in disorders related to the dopaminergic system.* . J Neural Transm Gen Sect. 1990;81(3):195–213. PubMed PMID: 2144427.
26. Messa C., Colombo C., Moresco R.M., Gobbo C., Galli L., Lucignani G., Gilardi M.C., Rizzo G., Smeraldi E., Zanardi R., Artigas F., Fazio F. *5-HT(2A) receptor binding is reduced in drug-naive and unchanged in SSRI-responder depressed patients compared to healthy controls: a PET study.* . Psychopharmacology (Berl). 2003;167(1):72–8. PubMed PMID: 12632246.
27. Moresco R.M., Colombo C., Fazio F., Bonfanti A., Lucignani G., Messa C., Gobbo C., Galli L., Del Sole A., Lucca A., Smeraldi E. *Effects of fluvoxamine treatment on the in vivo binding of [F-18]FESP in drug naive depressed patients: a PET study.* . Neuroimage. 2000;12(4):452–65. PubMed PMID: 10988039.
28. Lucignani G., Losa M., Moresco R.M., Del Sole A., Matarrese M., Bettinardi V., Mortini P., Giovanelli M., Fazio F. *Differentiation of clinically non-functioning pituitary adenomas from meningiomas and craniopharyngiomas by positron emission tomography with [18F]fluoro-ethyl-spiperone.* . Eur J Nucl Med. 1997;24(9):1149–55. PubMed PMID: 9283109.